
2016

Philadelphia Region of Pennsylvania

LGBTQA Community Health Needs Assessment Results





Study Purpose

This study assessed the prevalence of tobacco use, other chronic disease risk behaviors, and determinants of health in the lesbian, gay, bisexual, transgender, queer, and asexual (LGBTQA) communities in the Philadelphia region of Pennsylvania.

Methodology

Brief, anonymous, Internet-based surveys were completed by LGBTQA people in in the Philadelphia region of Pennsylvania, which included Bucks, Chester, Delaware, Montgomery, and Philadelphia counties.

During a 9-week period from March through May 2016, LGBTQA participants were purposively sampled using both direct and indirect recruitment strategies. Direct recruitment strategies included sending personal emails and Facebook messages to LGBTQA people. Indirect recruitment strategies included geographically targeted and LGBTQA-targeted Facebook advertisements, as well as email blasts.

Participants were informed that the data they provided were being collected anonymously and that they could stop the survey or refuse to answer any questions at any time. At the conclusion of the survey, participants were given the option to be redirected to an unlinked database where they could input their contact information to be entered into a lottery drawing for one \$50 gift card incentive.

Acknowledgements

This study was commissioned by the William Way LGBT Community Center with funding from the Pennsylvania Department of Health. Data were collected in partnership with the Mazzoni Center, Attic Youth Center, City of Brotherly Love Softball League, City of Philadelphia Department of Public Health, The COLOURS Organization, LGBT Elder Initiative, Stonewall Sports, and LGBTea Dances. Robert W.S. Coulter, M.P.H., served as the Statistical Data Analyst for this project: he conducted the data analyses and wrote the results. Scout, Ph.D., from LGBT Health Link, designed the questionnaire, helped guide survey administration, and drafted the recommendations for this report. Adrian Shanker from Bradbury-Sullivan LGBT Community Center provided technical assistance throughout the project.

Highlights

Sociodemographic Characteristics

In total, 1,218 LGBTQA individuals completed this survey. Nearly three-quarters (72.2%) of the sample identified as gay/lesbian, 10.5% as bisexual, and 14.4% as queer or pansexual. Additionally, 15.3% of the participants were transgender, 53.4% were cisgender men, and 30.3% were cisgender women.

Health Care Access & Quality

Health care coverage was highly prevalent in the sample (95.2%), and slightly higher than the prevalence of health care coverage among Pennsylvania adults (which was 89.9% according to the 2014 Behavioral Risk Factor Surveillance System [BRFSS]). About one-third (31.6%) of the sample had a health care provider react poorly to their LGBTQA status—and this varied by gender: transgender people (53.9%) were significantly more likely than cisgender women (31.9%) or men (25.2%) to have a provider react poorly. Additionally, transgender people (31.9%) were more likely than cisgender women (9.9%) and men (6.5%) to always or often fear a negative reaction by a health care provider. Nevertheless, most participants (89.0%) were out as LGBTQA to one or more of their health care providers.

LGBTQA Acceptance

LGBTQA participants thought their friends were the most LGBTQA-accepting (85.7% thought their friends were very LGBTQA-friendly). Second most accepting were participants' workplaces (51.9% thought their workplaces were very LGBTQA-friendly), followed by schools and birth families (40.7% and 40.5%, respectively, thought these groups were very LGBTQA-friendly). Medical providers and neighborhoods were not terribly accepting overall (35.3% and 29.0%, respectively, thought these groups were very LGBTQA-friendly).

Overall Health

In total, 69.2% of participants reported that their overall health status was good or very good. Importantly, participants' self-reported overall health status was significantly greater if they reported greater overall LGBTQA acceptance across the domains specified above. Nearly all respondents were interested in incorporating more healthy living strategies into their lives.

Body Mass Index

According to participants' body mass index (BMI), 33.7% were obese and 33.6% were overweight; this did not significantly vary by gender. Among the general population of Pennsylvania adults, 30.2% of adults were obese and 33.9% were overweight (BRFSS, 2014), suggesting that the LGBTQA population in the Philadelphia Region may be slightly more obese than the general Pennsylvanian population. Importantly, about two-thirds of LGBTQA participants were interested in health eating (63.2%) and active living strategies (67.6%).

Mental Health Treatment

In the past 12 months, 41.6% of participants received psychological counseling or treatment. In their lifetime, nearly two-thirds of the sample (63.7%) had taken medicine or received treatment for mental health problems.

HIV among Cisgender Men and Transgender People

HIV prevalence was significantly higher for cisgender men (12.5%) than transgender people (1.6%). Regular HIV testing was low. Excluding HIV-positive participants, 52.4% of cisgender men and 52.5% of transgender people received their last HIV test over one year ago. The Centers for Disease Control and Prevention (CDC) recommends that at-risk groups test for HIV at least once per year, and more often for those who engage in high-risk behaviors.

Tobacco Use & Opinions

In the past year, 34.3% of participants used any tobacco/nicotine products, and this significantly varied by gender: transgender people (46.2%) reported higher tobacco use than cisgender women (35.9%) and men (30.2%). Specifically, cigarette smoking was higher among LGBTQA participants than the general Pennsylvanian population (26.9% versus 19.9%, respectively; BRFSS, 2014). On a positive note, LGBTQA participants used smokeless tobacco (e.g., chew or snuff) less than the general Pennsylvanian population (1.1% vs. 4.3%; BRFSS, 2014).

LGBTQA people had little knowledge about smoking disparities: 37.4% strongly or somewhat agreed that LGBTQA people smoke more than the general population. Among all LGBTQA participants 83.4% were strongly or somewhat unappreciative that the tobacco industry advertises to the LGBTQA community; and 50.6% strongly or somewhat agreed that pride celebrations should be smoke-free events.

LGBTQA smokers overwhelmingly wanted to quit (89.8%), with the largest group of those wanting to quit within the next six months. When asked how they would quit, willingness to use effective cessation techniques was low: 32.0% of smokers would go to their health care provider, and 15.4% would enroll in a smoking cessation class or program; however, only 10.2% would use the quitline, and 17.7% of smokers did not know where to go for cessation assistance.

Cancer

The lifetime prevalence of cancer was 10.4% among LGBTQA participants (and this did not significantly vary by gender). Nearly one-third (31.2%) of participants who were assigned female at birth reported not being up to date with cervical Pap smears. Among participants 50 years or older: 27.4% were not up to date with colonoscopies; 14.6% of those assigned female at birth were not up to date with mammograms; and 17.6% of those assigned male at birth were not up to date with prostate exams. Overall, 88.7% of transgender people, 87.5% of cisgender women, and 77.7% of cisgender men never had an anal Pap smear. LGBTQA participants thought there was a high need for a variety of LGBTQA-welcoming practices for cancer care, including survivor support groups, caregiver support groups, inclusive hospital policies, and legal planning at hospitals.

Perceptions of LGBTQA Community Health Issues

Mental health, STDs/HIV, and suicide were perceived to be the top three health problems by LGBTQA participants.

Recommendations

Because tobacco use greatly affects the health of LGBTQA people, the lack of knowledge about LGBTQA smoking disparities and tobacco cessation programs are of particular concern. The following steps would help address this problem:

- Create LGBTQA-tailored smoking cessation groups;
- Train quitline providers to be LGBTQA friendly and affirming;
- Create LGBTQA-tailored quitline promotion campaigns to raise awareness for this cessation resource; and
- Create LGBTQA-tailored tobacco awareness campaigns to raise the knowledge level about LGBTQA smoking disparities and help people make healthier decisions.

Because many LGBTQA participants experienced LGBTQA-unfriendliness by a medical provider and perceived a high-need for LGBTQA-related cancer programs, enhancing the LGBTQA-friendliness of medical environments would have many benefits. The following steps would help address this need:


- Train medical providers to be LGBTQA culturally competent and to provide LGBTQA competent health care; and
- Design and disseminate LGBTQA-tailored materials that help make medical environments more LGBTQA-friendly.

To monitor the health of LGBTQA participants in the Philadelphia Region, LGBTQA health surveillance data should be routinely collected.


Results

In total, 1,218 LGBTQA individuals completed the survey. In this report, transgender is used as an umbrella term for the following individuals:

- People who identified as a trans man/woman or trans masculine/feminine;
- People who identified as agender, bigender, demigirl, fluid, non-conforming, non-binary, genderqueer, or Two-Spirit; and
- People whose assigned sex at birth does not match their current gender identity (e.g., someone who was assigned male sex at birth and now identifies as female).














How would you describe your neighborhood?
























































COMMUNITY



















































































If you smoke, what are your intentions to quit?





If you smoke, where would you most likely go for assistance to quit smoking?





unwelcoming

